

The forget-remember mechanism for 2-state spreading

J. Gu* and X. Cai†

*Complexity Science Center and Institute of Particle Physics,
Central China Normal University, Wuhan 430079, China*

(Dated: February 8, 2008)

A new mechanism, the forget-remember mechanism, is proposed for studying the spreading process in 2-state model. Such mechanism exhibits behaviors of message spreading influenced by some kinds of functions about time and history caring about the individuals of the spreading system, holding message or being out of message. To demonstrate the mechanism, both linear and exponential forms for forget-function and remember-function are simulated and show that a great impact on the saturation of message-spreading and the relative phase transformation.

PACS numbers: 87.23.Ge, 89.70.+c, 87.23.Kg

In this Letter we suggest a new mechanism, the mechanism with forget and remember functions, to study the message spreading in 2-state model and show their behaviors impressed on spreading process. Recent researches has widely addressed in the epidemic modelling[1, 2, 3, 4, 5, 6, 7] and the biological interaction, which has been shown valuable insights on spreading[8]. Various epidemiological models has been attracted attention of epidemiologists[9] and it was pointed out that epidemiological processes can be regarded as percolation[5, 10, 11, 12].

As known, most of spreading systems should be considered to have the following stronger resemblances: (i) In any of the spreading systems, there are certainly some special objects to be transmitted, which could be classified into two categories, things and messages. Things are not like messages. Things in a spreading system should be conservative, e.g. passengers on a flight were transported on airport network from City-A to arrive City-B, they are no longer in City A[13, 14]. But messages can be copied with some freedom and without conservation of the total number, e.g. a news you delivered to your friend, you still hold the news. (ii) Messages not only might be spread or lost, but also might be forgotten(not completely lost) or remember(after forgotten). (iii) Messages could be multifarious, such as virus on internet, e-mail, rumors in public, epidemic forest fire and so on[9, 15]. (iv) Each individual inside a message-spreading system could be at one of two states(holding message or being out of message) in 2-state models. And there still exist two cases, in the first case, an individual could transmit message to others or not, and in the second case, an individual could get message infected from the others or not. Altogether four states. No more other states for message-spreading systems. For example, the SI model assumes that each individual must be in one of two possible states, either susceptible(without message and can be transmitted in) or infective(with message

and can be transmitted out). The SIR model adds the third state[16, 17], removed state(lost message but can not be accepted message again). (v) Individual copies message into or from its neighbors, topologically only, not regarding the spacial distance and the limited transmitting speed between individuals usually. (vi) Time of evolution of the spreading system is sometimes considered discrete replacing continuous.

Based on the above-mentioned understanding which we prefer to call it the message theory to information theory, we will focus our attention on the influence of a mechanism, the forget-remember mechanism(FRM), in our investigation on message spreading. As we know, in the most of ready-made spreading studies, the ordinary means assumed a constant to measure the probability of spreading, and the most important parameter expressing the effective spreading rate, which not only to decide the percent of individuals with message, but also to determine whether the prevalence can break out[16, 17]. It is not a proper supposition here. Message in spreading maybe resemble knowledge in learning. In nature, the learning curve discovered by German Experimental Psychologist Hermann Ebbinghaus is famous[18, 19], which could be corresponding in spreading dynamics. Our so-called FRM is described now as following. Firstly, when an individual holds message, it may lose it with probability $P_{-}(t)$, a function on time t . We ponder a circumstance that the longer the individual holding message, the easier it forget message. Secondly, when an individual had seized message but lose it later, it can recall that message easier after shorter time with probability $P_{+}(t)$, also a function on time t . Taking notice, here the time in $P_{\pm}(t)$ in the FRM is not same to the system-time, but only cares about the history of each individual. The time in forget mechanism is zero-counting at the moment when the individual got message, while in remember mechanism, zero-counting at the moment when it lost message. We will show the influence of the FRM on time-interval and how it to impact on spreading curve.

We attempt to consider our FRM by using a developed model, e.g. the Barabási-Albert(BA) algorithm as a frame for message-spreading, because it can reflect the social network strikingly and its degree distribution ex-

*Electronic address: guj@iopp.cnu.edu.cn

†Electronic address: xcai@mail.cnu.edu.cn




FIG. 1: Functions of the FRM: the linear form and the exponential form. $P_-(t)$ is probability for the forget mechanism while $P_+(t)$ for the remember mechanism, both are ranged from zero to one.

hibits a power-law behavior of with form $P(k) \sim k^{-\gamma}$ where exponent γ ranged between 2 and 3[15, 20]. In Ref.[9, 20, 21] the epidemic processes in an uncorrelated network do not possess an epidemic threshold, and below which the diseases cannot produce a macroscopic epidemic outbreak. In a spread process, if an infected individual transmits message with probability δ , and a susceptible individual is infected by its infected neighbor with probability ν , the proportion of ν and δ is defined as an effective spreading rate λ . Taking the BA networks, and let $\rho_k(t)$ the probability of a message-holding individual with degree k , we get an equation for the mean field function

$$\frac{\partial \rho_k(t)}{\partial t} = -\rho_k(t) + \lambda k [1 - \rho_k(t)] \Theta(\rho_k(t)) \quad (1)$$

where $\Theta(\rho_k(t))$ is the probability of any given link to a message-holding individual[9]. Then we can reach a spread threshold $\lambda_c = \langle k \rangle / \langle k^2 \rangle$, so that $\lambda_c \rightarrow 0$ if amount of individuals in a network $N \rightarrow \infty$ [22, 23].

Choosing the BA networks built with the preferential attachment character[15] with amount $N = 1000$. To beginning with an initial network with $m_0 = 5$ individuals, a new individual i is added to the network linked to an existing individual j with probability: $\Pi_i = k_i / \sum_j k_j$, where k_i and k_j are degrees of individuals i and j . The average degree $\langle k \rangle = 3.8$ and the exponent $\gamma = 2.2$. Now let us address the forget mechanism. We assume that an individual without message on the network can get message from its neighbors which are holding message with probability ν . An individual which is in a message-holding state may become into a message-losing state with the probability function $P_-(t)$, where concerning

the time t , i.e. how long this individual holding message. Certainly, $t > 0$ and the values of $P_-(t)$ range from zero to one. Meanwhile the remember mechanism is built by considering an individual in a message-losing state, but it can recall message only if it had possessed of this message before. This case is common in natural, especially it takes place on those processes, e.g. language-learning, disease-catching or rumors-circulating repeatedly and so on. An individual at a state without message, but which must formerly have gotten message by copying or infected from its neighbors, can remember message controlled by the probability function $P_+(t)$. Usually the longer the individual was out of the message, the harder it remember the message. The ability to recall the past should be concerning the history, that is to say, if an individual had never owned message and then had never undergone on a forget-process, it can not remember any message. Time t in $P_+(t)$ is zero-counting when it lost message for each individual respectively.

In order to gain further insight into the FRM, we consider two forms for both $P_-(t)$ and $P_+(t)$: the linear form and the exponential form, see Eq.(2) with parameters a and b and Eq.(3) with α and β . To make our mechanism clear, parameter b or β had better take more than zero to decide the shapes of probability functions for forget and remember mechanisms with additional constant terms a or α . By the way, when b or β is equal to zero, i.e. the probability functions should be simply constant, we shall discuss the first form only.




FIG. 2: Plot of the density $D(t)$ versus the system-time step t with the spreading probability $\nu = 0.002$. Figures on the left-row for forget mechanism and on the right-row for remember mechanism. The dashed curves are simulated with fixed remember probability $P_+ = 0.1$ while the solid curves without remember mechanism in left figures. Figures on the upper three lines by using linear form with values of parameters a and b , and those on the lower two lines by using exponential form with values of parameter α and β as shown.

The linear function is defined as

$$P_{\mp}(t) = \begin{cases} 0 & \text{for } \pm t < -a/b, \\ a \pm bt & \text{for } \pm t \in [-a/b, (1-a)/b], \\ 1 & \text{for } \pm t > (1-a)/b. \end{cases} \quad (2)$$

and the exponential function as

$$P_{\mp}(t) = \begin{cases} 0 & \text{for } \mp e^{-\beta t} > -\alpha, \\ \alpha \mp e^{-\beta t} & \text{for } \mp e^{-\beta t} \in [1-\alpha, -\alpha], \\ 1 & \text{for } \mp e^{-\beta t} < 1-\alpha. \end{cases} \quad (3)$$

Since $P_{\mp}(t)$ are probabilities for forget and remember functions, they are ranged between 0 and 1 as shown in Fig.1. The behavior of the influence of the above different forms to the spreading system can be computed by noticing that the zero-counting of time t at which each individual forgets or remembers message separately. let $D(t)$ be the density of individuals with message on the




FIG. 3: Saturation density D_s versus parameters in different forms. Figures on the left-row for forget mechanism and on the right-row for remember mechanism. Figures on the upper three lines by using the linear form with a and b , and those on the lower two lines by using the exponential form with α and β as shown.

frame in the BA model. As showed most obviously in Fig.2, the influences make a faster progress to the spreading process in the early time-steps of the evolution, and always reach saturation, i.e. not converting whole individuals of the system after long time spreading, both in linear and exponential forms.

Not matter how many the parameters were taken in both forms with the forget mechanism, the value of $D(t)$ in the saturation is descending with parameters increasing. And with the remember mechanism, since the value of function is ascending with parameters a and α increasing, the $D(t)$ at the saturated moment is also rising with them, but is opposite with the parameters b and β . As shown, in the left five figures of Fig.2, the dashed curves are considered the forget mechanism with fixed remember probability while the solid curves without remember mechanism. The results show that the density $D(t)$ in the case of the left figures might be easy saturated. The reason is that the remember mechanism inspires the prevalence of message, while the forget mechanism restrains the spreading.

Let D_s be the saturated value of $D(t)$ in different cases. Fig.3 exhibits the relation between D_s and parameters a , b and α , β . Noteworthy, as shown in Fig.3 the saturation density D_s can vary in a very large range (from 0 to 1) controlled by parameter taking. That is to say, the FRM is not a slimly adjustment on the spreading process, which is to the contrary to the conclusion that only the effective spreading rate is important to the spreading process in some fashionable models. In our figures, part of the relations between the saturation density D_s and parameters could be fitted by the exponential function. And as shown, there exists the thrill of a new sight which is no other than the phase transformation points about the saturation density D_s . At the phase transformation point, the data are divided into two sides, on the one side, D_s is ascending or descending quickly, and on the other side, D_s changes very slowly. This observation based on the FRM is different from the present study of the spreading process on the BA network[20, 21].

In summary, the FRM we have presented for spreading process with 2-state systems is possibly the simplest one in a kind of selected practical networks(the BA model) currently in effect. A novel feature in the model with our forget-remember mechanism is the saturation of the spreading density with phase transition controlled by the parameters, i.e. the forget coefficients and the remember coefficients. The results can be concluded as follow:

(i) The probability in spreading process should not be always constant, maybe depends on time t , especially cares about the history of each individual holding the message. (ii) Evidently the forget mechanism restrains the spreading but the remember mechanism inspires it. Truth should be simple and obvious. (iii) The effective spreading rate is not the only possible important factor to decide the density of the message-holding individuals and whether the prevalence can break out. The FRM would naturally work on the spreading systems to let the spreading process reach the saturation with varying value ranged widely from zero to one at most. It serves as a great source of inspiration for the spreading process control. (iv) There exists the phase transformation on the saturation of spreading density. It needs to investigate further.

The FRM on the spreading system can explain why different diseases have different saturation in population, even though they have the semblance effective spreading rate. Different attitudes derive from different perspectives. It is sometimes better to emphasize the attention to history of each individual inside the system. Details is the key of characteristic of the whole.

This work is supported by the National Natural Science Foundation of China (Grant Nos. 70571027, 10647125, 10635020 and 70401020) and the Ministry of Education of China(Grant No. 306022).

-
- [1] P. Echenique, J. Gomez-Gardeñes, and Y. Moreno, Phys. Rev. E **70**, 036102 (2004).
 - [2] L. Zhao, Y. C. Lai, K. Park, and N. Ye, Phys. Rev. E **71**, 026125 (2005).
 - [3] L. Donetti, P. I. Hurtado, and M. A. Muñoz, Phys. Rev. Lett. **95**, 188701 (2005).
 - [4] Z. Y. Chen and X. F. Wang, Phys. Rev. E **73**, 036107 (2006).
 - [5] M. E. J. Newman, Phys. Rev. Lett. **95**, 108701 (2005); Phys. Rev. E **65**, 021904 (2002).
 - [6] T. Gross, C. J. D. D'Lima, and B. Blasius, Phys. Rev. Lett. **96**, 208701 (2006).
 - [7] J. D. Noh and H. Park, Phys. Rev. Lett. **94**, 145702 (2005); Phys. Rev. Lett. **92**, 118701 (2004).
 - [8] O. Sporns, D. R. Chialvo, M. Kaiser and C. C. Ilgetag. TRENDS in Cognitive Sciences Vol.8, No.9, 418-425 (2004).
 - [9] R. Pastor-Satorras, A. Vázquez, and A. Vespignani, Phys. Rev. Lett. **86**, 14 (2001); Phys. Rev. E **65**, 035108 (2002); Phys. Rev. E **65**, 035108 (2006); Phys. Rev. E **63**, 066117 (2001).
 - [10] A. Jiménez-Dalmaroni, Phys. Rev. E **74**, 011123 (2006).
 - [11] N. Wagner, I. Balberg and D. Klein, Phys. Rev. E **74**, 011127 (2006).
 - [12] M. A. Serrano and M. Bogguá, Phys. Rev. E **74**, 056115 (2006).
 - [13] W. Li and X. Cai, Phys. Rev. E **69**, 046106 (2004).
 - [14] L. P. Chi, R. Wang, et al., Chin. Phys. Lett. **20**, No.8, 1393 (2003).
 - [15] A-L. Barabási and R. Albert, Science, **286**, 509 (1999); Rev. Mod. Phys. **74**, 47 (2002); Science, **286**(5439), 509-512 (1999).
 - [16] R. M. Anderson, R. M. May. Infections Diseases in Humans, Oxford University Press, Oxford, (1992).
 - [17] J. D. Murray, Mathematical Biology, Springer, Berlin, (1993).
 - [18] linked to <http://psychclassics.yorku.ca/Ebbinghaus/wozniak.htm>.
 - [19] Memory: A Contribution to Experimental Psychology Hermann Ebbinghaus (1885), translated by Henry A. Ruger & Clara E. Bussenius (1913), originally published in New York by Teachers College, Columbia Universit.
 - [20] A-L. Barabási, R. Albert and H. Jeong, Physica(Amsterdam) **272A**, 173 (1999).
 - [21] Robert M. May and Alun L. Lloyd, Phys. Rev. E **64**, 066112 (2001).
 - [22] Diekmann O, Heesterbeek J A P. Mathematical Epidemiology of Infectious Disease: Model Building, Analysis and Interpretation. New York: John Wiley & Son publisher, (2000).
 - [23] R. Pastor-Satorras1 and A. Vespignani, Epidemics and immunization in scale-free networks. In Handbook of Graphs and Networks, Bornholdt S., Schuster H. G.(eds), WILEY-YVCH publisher, (2003).